

321
CLAIMS 1-21

What is claimed is;

1. An immunostimulating peptide having an amino acid sequence X_1 LYQYMDDV, wherein X_1 is any hydrophobic amino acid.

2. The immunostimulating peptide of claim 1, wherein the amino acid sequence is VLYQYMDDV.

3. A medicament comprising:
 i) the immunostimulating peptide of claim 1; and,
 ii) a pharmaceutically acceptable excipient.

4. The medicament of claim 3, further comprising an immunostimulant.

5. A method for preventing or treating an HIV-1 infection comprising administering a dose of the medicament of claim 3 in an amount effective to induce an immune response capable of preventing HIV-1 infection or reducing HIV-1 viral load in a patient.

6. The method of claim 5, wherein the patient is a human.

7. A immunostimulating peptide or protein comprising the sequence X_1X_2 YQYMDDV X_3 wherein

X_1 is a sequence of amino acid residues of between 0 and 200 residues in length;

X_2 is any hydrophobic amino acid; and,

X_3 is a sequence of amino acid residues of between 0 and 200 residues in length.

8. A medicament comprising a vector including a nucleic acid comprising a nucleotide sequence encoding a peptide having the sequence X_1 LYQYMDDV, wherein X_1 is any hydrophobic amino acid,
 wherein introducing the medicament to a subject results in

expression of the nucleic acid, thereby inducing an immune response in the subject directed against an epitope of a product encoded by the nucleic acid.

9. The medicament of claim 8, wherein the vector is a virus.

10. A method for preventing or treating an HIV-1 infection comprising administering a dose of the medicament of claim 8 in an amount effective to induce an immune response capable of preventing HIV-1 infection or reducing HIV-1 viral load in a patient.

11. A method of ⁽¹⁾ assessing ⁽²⁾ immune function or diagnosing exposure to HIV-1 for a subject, the method comprising:

- i) contacting a blood sample comprising T cells obtained from the subject with a peptide having an amino acid sequence X₁LYQYMDDV, wherein X₁ is any hydrophobic amino acid; and,
- ii) determining an immune response of the subject's T cells to the peptide. ?

12. The method of claim 11, wherein said determining step (ii) is performed by assaying for IFN-γ production, or lysis of cells displaying the peptide by cytotoxic T lymphocytes induced with the peptide.

13. A fusion molecule comprising an amino acid sequence X₁LYQYMDDV, wherein X₁ is any hydrophobic amino acid.

14. The fusion molecule of claim 13, further comprising an amino acid sequence for an HIV-1 viral protein

15. The fusion molecule of claim 13, further comprising a glycolipid.

16. The fusion molecule of claim 13, further comprising an amino acid sequence for an immunostimulating carrier protein.

17. A peptide or protein comprising an amino acid sequence X₁LYQYMDDV, wherein X₁ is any hydrophobic amino acid.

18. The peptide or protein of claim 17, further comprising an acetylated N-terminus.

19. The peptide or protein of claim 17, further comprising a modification to the C-terminus, the modification selected from the group consisting of amidation, esterification, and reduction of a C-terminal amino acid carboxyl group..

20. A medicament comprising a peptide of claim 1 pulsed onto dendritic cells.

21. A medicament comprising dendritic cells transduced with a vector of claim 8